Western Washington Poisonous Plants

Western Washington Plants Known to Be Poisonous to Horses (USDA-SCS)

Buttercup, **creeping** (Ranunculus repens)

Location or Season	Moist soils
Probable Toxic Dose	Very Large
Toxin	Protoaneonin
Symptoms	Inflammation and narcosis
Comments	Rarely eaten unless pasture overgrazed

Camas, death (Zigadenus venenosus)

Location or Season	Spring
Probable Toxic Dose	less than 9 lb.
Toxin	Steroidal, glycosidal, alkaloid
Symptoms	Salivation, weakness, respiratory difficulty, nausea, convulsions, coma
Comments	Deadly, easily confused with edible camas after bloom

Fern, bracken (Pteridium aquilinum)

· · · · · · · · · · · · · · · · · · ·	
Location or Season	Fall, when pastures overgrazed
Probable Toxic Dose	Cumulative Large quantities
Toxin	Thiaminase
Symptoms	Appetite loss, timid, stupified, incoordination
Comments	

Fiddleneck (Amsinckia sp.)

The state of the s	
Location or Season	Overgrazed pastures
Probable Toxic Dose	Single dose 20 lb. or cumulative
Toxin	pyrrolixidine
Symptoms	Liver damage & failure, depression, dermatitis, incoordination, death
Comments	Similar poison to Tansy Ragwort

Foxglove (Digitalis purpurea)

ongrove (Bightans parparea)	
Location or Season	Acid soils
Probable Toxic Dose	VERY TOXIC (1/4 lb.)
Toxin	Digitoxin & other glycosides
Symptoms	Contracted pupils, labored breathing, convulsions, death
Comments	Rarely eaten fresh, dangerous in hay

Hemlock, poison (Conium maculatum)

remireday person (comunication	,
Location or Season	Ditches, moist disturbed areas
Probable Toxic Dose	Very toxic (5 to 10 lb.)
Toxin	Coniine and other alkaloids
Symptoms	Narcosis, paralysis, death
Comments	Hay MAY be somewhat less toxic as the poison will slowly evaporate

Hemlock, water (Cicuta douglasii)

Location or Season	Low, wet areas; in spring roots pull out of ground easily
Probable Toxic Dose	VERY TOXIC (.2 to .8 lb.)
Toxin	Circutoxin
Symptoms	Teeth grinding, muscle spasms, respiratory failure, death
Comments	Roots & stem base most toxic. The most poisonous plant known in North America

Horsetail (Equisetum arvense)

Horsetan (Equisetani di Vense)	
Location or Season	Moist areas
Probable Toxic Dose	Large quantities Cumulative
Toxin	Thiaminase
Symptoms	Thiamine deficiency causes appetite loss, incoordination
Comments	Poisoning occurs when dry plants are fed in hay

Knapweed, Russian & Yellowstar thistle (Centaurca spp.)

triapriced, itassian e renovistar tristic (centaured spp.)	
Location or Season	Disturbed areas
Probable Toxic Dose	Cumulative 600 lb.?
Toxin	
Symptoms	Brain deterioration resulting in "Chewing disease"
Comments	Can eventually cause death by starvation<

Larkspur (Delphinium spp.)

Eurkspur (Deiphinnum spp.)	
Location or Season	
Probable Toxic Dose	Very toxic (1 lb.)
Toxin	Alkaloids-delphinine
Symptoms	Constipation, bloat, depression, paralysis
Comments	Deadly

Western Washington Poisonous Plants

Western Washington Plants Known to Be Poisonous to Horses (USDA-SCS)

Lupine (Lupinus spp.)

-upc (-upus spp.)	
Location or Season	
Probable Toxic Dose	600 to 800 lbs.
Toxin	Alkaloids-lupinine
Symptoms	Spasms, cerebral excitement & death
Comments	Can cause birth defects if eaten during pregnancy

Nightshade, black (Solanum spp.)

<u> </u>	11 /
Location or Season	Late summer, early fall, fencerows
Probable Toxic Dose	1 to 10 lb.
Toxin	Alkaloid-solanine
Symptoms	Diarrhea, convulsions, incoordination, death
Comments	

Oak (Ouercus spp.)

Location or Season	
Probable Toxic Dose	Very large
Toxin	Tannins?
Symptoms	Constipation, blood in urine
Comments	Leaves can cause problems, acorn poisoning more common

Ragwort, tansy & Common groundsel (Seneclo spn.)

rtugirort, turisy & common g	grounder (serieus spp.)
Location or Season	
Probable Toxic Dose	
Toxin	Alkaloid-pyrrolizidine
Symptoms	Liver Lesions, weakness, staggering, death
Comments	Liver damage is permanent. Normally avoided when fresh, eaten in hay or when wilted.

Rhododendron (Rhododendron spp.)

	1
Location or Season	
Probable Toxic Dose	Small
Toxin	Glycosides
Symptoms	Vomiting, vertigo, death from respiratory failure
Comments	

Yew (Taxus spp.)

Location or Season	
Probable Toxic Dose	1 to 10 lbs.
Toxin	Alkaloid-taxine
Symptoms	Gastroenteritis, labored breathing, trembling, collapse
Comments	Rarely eaten fresh, dangerous in hay

Excerpted from: <u>Pasture Management for Horses and Ponies</u>, Gillian McCarthy; and *Plants that Poison Livestock in Thurston County*, Thurston County Noxious Weed Control Agency.

Other resources:

University of Illinois, Urbana-Champagne

http://www.library.uiuc.edu/vex/toxic/intro.htm http://www.library.uiuc.edu/vex/toxic/comlist.htm

Cornell University

http://www.ansci.cornell.edu/plants/plants.html http://www.ansci.cornell.edu/toxhorses.html

Purdue University

http://vet.purdue.edu/depts/addl/toxic/cover1.htm http://vet.purdue.edu/depts/addl/toxic/byanim.htm

HorseAdvice.com

http://www.horseadvice.com/articles/diseases/poisonmenu.html

in Canada

http://res.agr.ca/brd/poisonpl/poisons.html